


One more question dealing with horizontally launched projectiles.

Ex) Will a tennis ball, served horizontally at 40.0 m/s, from a height of 2.2 m clear a net 0.90 m high and 10.0 m away?



$$t = \frac{d_x}{v_x} = \frac{10 \text{ m}}{40 \text{ m/s}} = 0.25 \text{ s}$$

$$d_y = v_{iy}t + \frac{1}{2}at^2$$

$$d_y = \frac{1}{2}(-9.8 \text{ m/s}^2)(0.25 \text{ s})^2$$

$$d_y = -0.31 \text{ m}$$

This means the ball fell 0.31 m from its initial height.

$$d = 2.2 \text{ m} - 0.31 \text{ m}$$

$$d = 1.89 \text{ m or } 1.9 \text{ m}$$

After 0.25 s the ball is 1.9 m above the ground, so it will clear a net 0.9 m high.

Section 1.4 Launching Projectiles at Any Angle


In this lesson

You will calculate:

- the total time of the motion when the initial velocity and initial height are given
- the range and final velocity when the initial velocity and initial height are given
- the velocity and displacement at anytime when the initial velocity and initial height are given
- the height from which the projectile is fired when its final velocity is given
(e.g., the projectile may be fired at an angle from a roof top or from a cliff.)
- the initial velocity when the final velocity and change in position are given

A projectile is launched at an angle above the horizontal as shown below.


0
3
4
5



- v_1 is the **initial velocity** of the ball. It is directed at an angle θ above the horizontal. The initial **vertical** component of the velocity is v_{1y} and the initial **horizontal** component of the velocity is v_{1x} .
- The vertical velocity decreases until it becomes 0.0 m/s at the very top of the trajectory. Then, it increases in exactly the same increments as the ball falls back to earth. **Just as the ball reaches the level at which it started, the magnitudes of v_{1y} and v_{2y} are identical.** These two velocities are, of course, in exactly opposite directions.
- Assuming no air friction (even though that's unreal!) there is no force either aiding or opposing the horizontal vectors. Therefore the **horizontal components, v_x , remain constant.**
- Assuming the ball returns to the level at which it started, the magnitudes of v_1 and v_2 are the same. The difference is that v_1 makes an angle of θ above the horizontal, while v_2 makes an angle θ below the horizontal.

Launch Angles

Suppose a cannon ball is launched at a speed of 20.0 m/s at 15°, 30°, 45°, 60°, 75°.



15°	$v_x = 20 \frac{m}{s} \cos 15^\circ$	$v_y = 20 \frac{m}{s} \sin 15^\circ$
	$= 19.3 \frac{m}{s}$	$= 5.2 \frac{m}{s}$
30°	$v_x = 20 \frac{m}{s} \cos 30^\circ$	$v_y = 20 \frac{m}{s} \sin 30^\circ$
	$= 17.3 \frac{m}{s}$	$= 10 \frac{m}{s}$
45°	$v_x = 14.1 \frac{m}{s}$	$v_y = 14.1 \frac{m}{s}$
60°	$v_x = 10 \frac{m}{s}$	$v_y = 17.3 \frac{m}{s}$
75°	$v_x = 5.2 \frac{m}{s}$	$v_y = 19.3 \frac{m}{s}$

1 Which ball is in the air for the shortest period of time?

The one launched at 15° will hit the ground first. The **hang time** of a projectile (i.e., the length of time the ball is in the air) is determined by the initial value of the vertical velocity component (v_{1y}). The smaller v_{1y} , the shorter the hang time.

2 Which ball has the highest peak?

The one launched at 75° . The peak height is determined by v_{1y} . The greater v_{1y} , the higher it will rise.


3 Which ball has the greatest range?

The one launched at 45° . Range is determined by v_{1x} and the hang time is determined by v_{1y} .

Examples

- 1 A soccer ball is kicked so that its initial velocity is 18.2 m/s at an angle of 54° with the field.

A How long is the ball in the air?



$$\begin{aligned} \frac{H}{V_{ix}} &= 18.2 \text{ m/s} \cos 54^\circ \\ &= 10.7 \text{ m/s} \end{aligned}$$

$$t = \frac{V_{2y} - V_{1y}}{a}$$

$$t = \frac{-14.7 \text{ m/s} - 14.7 \text{ m/s}}{-9.8 \text{ m/s}^2}$$

$$t = 3.0 \text{ s}$$

$$\begin{aligned} \frac{V}{V_{iy}} &= 18.2 \text{ m/s} \sin 54^\circ \\ &= 14.7 \text{ m/s} \\ a &= -9.8 \text{ m/s}^2 \\ V_{2y} &= -14.7 \text{ m/s} \\ &\text{(when it hits ground)} \end{aligned}$$

B Determine the range of the ball.

$$\begin{aligned} dx &= V_x t \\ &= (10.7 \text{ m/s})(3.0 \text{ s}) \\ &= 32 \text{ m} \end{aligned}$$

- C Determine the speed and direction with which the ball hits the ground.

$$v_2 = 18 \text{ m/s at } 54^\circ \text{ below the horizontal.}$$


v_1 & v_2 are at ground level so v_1 & v_2 are equal in magnitude but opposite in direction.

- D How high did the ball rise above the field? * use 1.5s not 3.0s

$$\begin{aligned} d &= v_1 t + \frac{1}{2} a t^2 \\ &= (14.7 \text{ m/s})(1.5 \text{ s}) + \frac{1}{2} (-9.8 \text{ m/s}^2)(1.5 \text{ s})^2 \\ &= 11 \text{ m} \end{aligned}$$

- 2 A ball is kicked so that its initial velocity is 18.2 m/s at an angle of 54° with the playing field. This time the ball cleared an embankment and landed on a sandy beach 3.4 m below the level of the playing field.

- A How fast was the ball traveling when it hit the beach and what was its direction?



$$v_{2y}^2 = v_{1y}^2 + 2ad$$

$$= (14.7 \text{ m/s})^2 + 2(-9.8 \text{ m/s}^2)(-3.4 \text{ m})$$

$$v_{2y} = -16.8 \text{ m/s}$$

$$v_2^2 = v_{2x}^2 + v_{2y}^2$$

$$= (10.7 \text{ m/s})^2 + (16.8 \text{ m/s})^2$$

$$v_2 = 19.9 \text{ m/s at } 58^\circ \text{ below horizontal.}$$

$$\tan \theta = \frac{v_y}{v_x}$$

$$= \frac{16.8 \text{ m/s}}{10.7 \text{ m/s}}$$

$$\theta = 58^\circ$$

- B How much longer was the ball in the air (compared to the ball in question 1)?

$$t = \frac{v_{2y} - v_{1y}}{a}$$


$$= \frac{-16.8 \text{ m/s} - 14.7 \text{ m/s}}{-9.8 \text{ m/s}^2}$$

$$= 3.2 \text{ s}$$

The ball was in the air an extra 0.20 s. ($3.2 \text{ s} - 3.0 \text{ s} = 0.2 \text{ s}$)

- 3 A novice tennis player lobs the ball in a high arc as shown. A radar device clocks the ball at 10.9 m/s as it hits the court 1.83 s after being struck by the racket. The player has a reach of 0.75 m.

A How tall is the player?



H

$$V_{2x} = 10.9 \text{ m/s} \cos 70^\circ$$

$$= 3.73 \text{ m/s}$$

$$t = 1.83 \text{ s}$$

V

$$V_{2y} = 10.9 \text{ m/s} \sin 70^\circ$$

$$= 10.2 \text{ m/s}$$

use $V_{2y} = -10.2 \text{ m/s}$

$$a = -9.8 \text{ m/s}^2$$

$$t = 1.83 \text{ s}$$

$$d = V_2 t - \frac{1}{2} a t^2$$

$$d = (-10.2 \frac{\text{m}}{\text{s}})(1.83 \text{ s}) - \frac{1}{2}(-9.8 \frac{\text{m}}{\text{s}^2})(1.83 \text{ s})^2$$

$$d = -2.26 \text{ m}$$

∴ height ball was hit is 2.26 m

$$\text{height of player} = 2.26 \text{ m} - 0.75 \text{ m}$$

$$= 1.51 \text{ m}$$

B How fast was the tennis ball traveling when first struck by the racket?

($V_1 = ?$) Find V_{1x} & V_{1y}

$$V_{1x} = 3.73 \text{ m/s}$$

$$V_{1y} = V_{2y} - at$$

$$= -10.2 \text{ m/s} - (-9.8 \text{ m/s}^2)(1.83 \text{ s})$$

$$= 7.73 \text{ m/s}$$


$$V_1^2 = (3.73 \text{ m/s})^2 + (7.73 \text{ m/s})^2$$

$$V_1 = 8.6 \text{ m/s} \text{ at } 64^\circ \text{ above the horizontal}$$

$$\tan \theta = \frac{7.73 \text{ m/s}}{3.73 \text{ m/s}}$$

- 4 A ball is shot out of a cannon with a horizontal velocity component of 40.0 m/s and a vertical velocity component of 20.0 m/s. The cannon is sitting at the top of a 100.0 m high cliff.

A How far will it travel? (range)



$$v_{2y}^2 = v_{1y}^2 + 2ad$$

$$= (20 \text{ m/s})^2 + 2(-9.8 \text{ m/s}^2)(-100 \text{ m})$$

$$v_{2y} = \pm 48.6 \text{ m/s}$$

$$v_{2y} = -48.6 \text{ m/s}$$

$$t = \frac{v_{2y} - v_{1y}}{a} = \frac{-48.6 \text{ m/s} - 20 \text{ m/s}}{-9.8 \text{ m/s}^2}$$

$$t = 7.0 \text{ s}$$

$$dx = v_x t = (40 \text{ m/s})(7.0 \text{ s})$$

$$= 280 \text{ m}$$

B What will be the cannon ball's maximum height?

$$v_{1y} = 20 \text{ m/s}$$

$$a = -9.8 \text{ m/s}^2$$

$$v_{2y} = 0$$

$$d = ?$$

$$dy = \frac{v_{2y}^2 - v_{1y}^2}{2a}$$

$$dy = \frac{0 - (20 \text{ m/s})^2}{2(-9.8 \text{ m/s}^2)}$$

$$dy = 20.4 \text{ m}$$

wrt cliff: max $h = 20.4 \text{ m}$

wrt ground max $h = 120.4 \text{ m}$ or $1.20 \times 10^2 \text{ m}$

Homework: 19, 29 - 31, 33 page 115 - 116